

WEIGHTED PLB-SPACES OF CONTINUOUS FUNCTIONS ARISING AS TENSOR PRODUCTS OF A FRÉCHET AND A DF-SPACE

SVEN-AKE WEGNER ^a

ABSTRACT. Countable projective limits of countable inductive limits, so-called PLB-spaces, of weighted Banach spaces of continuous functions have recently been investigated by Agethen, Bierstedt and Bonet, who analyzed locally convex properties in terms of the defining double sequence of weights. We complement their results by considering a defining sequence which is the product of two single sequences. By associating these two sequences with a weighted Fréchet, resp. LB-space of continuous functions or with two weighted Fréchet spaces (by taking the reciprocal of one of the sequences) we derive a representation of the PLB-space as the tensor product of a Fréchet and a DF-space and exhibit a connection between the invariants (DN) and (Ω) for Fréchet spaces and locally convex properties of the PLB-space resp. of the forementioned tensor product.

1. INTRODUCTION

Agethen, Bierstedt, Bonet [2] investigated the structure of spaces of continuous functions defined on a locally compact and σ -compact topological space that can be written as a countable intersection of countable unions of weighted Banach spaces of continuous functions. These spaces were introduced and studied for the first time in Agethen's thesis [1]; they are examples of so-called PLB-spaces, i.e. countable projective limits of countable inductive limits of Banach spaces. Spaces of this type arise naturally in analysis, for instance the space of distributions, the space of real analytic functions and several spaces of ultradistributions are of this type. In fact, all the forementioned spaces are even PLS-spaces, that is the linking maps in the inductive spectra of Banach spaces are compact and some of them even appear to be PLN-spaces (i.e. the linking maps are nuclear). During the last years the theory of PLS-spaces has played an important role in the application of abstract functional analytic methods to several classical problems in analysis. In particular there was an intense research on tensor products of PLS-spaces motivated by the problem of parameter dependence of solutions of PDE, see e.g. Bonet, Domański [10, 12], Domański [15] and Piszczek [26]. We refer to the survey article [14] of Domański for applications, examples and further references.

Many of the applications reviewed by Domański [14] are based on the theory of the so-called first derived functor of the projective limit functor. This method has its origin in the application of homological algebra to functional analysis. The research on this subject was started by Palamodov [23, 24] in the late sixties and carried on since the mid eighties by Vogt [31] and many others. We refer to the book of Wengenroth [38], who laid down a systematic study of homological tools in functional analysis and in particular presents many ready-for-use results concerning concrete analytic problems. In particular, [38, section 5] illustrates that for the splitting theory of Fréchet or more general locally convex spaces, the consideration of PLB-spaces which are not PLS-spaces is indispensable.

Agethen, Bierstedt, Bonet [2] described locally convex properties (i.e. ultrabornologicity and barrelledness) of weighted PLB-spaces of continuous functions in terms of the defining double sequence of weights by using the theory of the derived projective limit functor Proj^1 , which we mentioned already above. In addition they studied the interchangeability of projective and inductive limit, i.e. the question when the PLB-spaces are equal to the weighted LF-spaces of continuous functions introduced by Bierstedt, Bonet [6]. Moreover, the work of Agethen, Bierstedt, Bonet exhibits that certain spaces of linear and continuous operators between Köthe echelon spaces as well as certain tensor products of a Köthe echelon and a co-echelon space happen to be weighted PLB-spaces of continuous functions, see [2, section 4].

In this paper we complement the results of [2] by considering the following special situation. We assume that the domain of the functions in the PLB-space is the product of two topological spaces and that the

2010 *Mathematics Subject Classification*: Primary 46E10; Secondary 46A13, 46M40, 46M05.

Key words and phrases: PLB-space, weighted spaces of continuous functions, tensor product of a Fréchet and a DF-space.

Date: October 10, 2010.

^aFB C – Mathematics, University of Wuppertal, Gaußstr. 20, 42119 Wuppertal, Germany, Phone: +49 202 439 253 1, Fax: +49 202 439 372 4, eMail: wegner@math.uni-wuppertal.de.

defining double sequence of weights is the product of an increasing and a decreasing (single) sequence on each of the two topological spaces. By taking the reciprocal of the decreasing sequence we can associate the double sequence with two weighted Fréchet spaces of continuous functions. This enables us to exhibit a connection between locally convex properties of the PLB-space and the invariants (DN) and (Ω) for Fréchet spaces. The latter were introduced by Vogt [29] and Vogt, Wagner [35] and play a prominent role in the structure theory of Fréchet spaces, see for instance the book [22] of Meise, Vogt. In particular we establish a criterion for ultrabornologicity of the PLB-spaces formulated in terms of (DN) and (Ω) . Taking no reciprocals the two sequences give rise to a weighted Fréchet and a weighted LB-space of continuous functions. In analogy to the case of sequence spaces (cf. [2, section 4]) we show that the ε -tensor product of these two spaces is isomorphic to the PLB-space defined by the double sequence explained above if we assume the decreasing sequence to be regularly decreasing (see Bierstedt, Meise, Summers [8, Definition 2.1]). Combining these results we finally obtain a criterion for the ultrabornologicity of the tensor product of a weighted Fréchet space of continuous functions and a weighted LB-space of continuous functions.

The underlying general question of determining topological properties of the tensor product of a Fréchet space and a DF-space was raised by Grothendieck in the last section of his thèse [17]. He investigated the case of π -tensor products of echelon and coechelon spaces of order one, see [17, Chapitre II, §4, No. 3, Theorem 15]. His results inspired many authors to further studies; see Varol [28, Section 0] for references and a generalization of Grothendieck's original result to the case of π -tensor products of a Köthe coechelon space of order one and arbitrary Fréchet spaces ([28, Theorem 2.1]). Varol [28, Theorem 2.7] investigated in addition the case of ε -tensor products of Köthe coechelon spaces of order zero and arbitrary Fréchet spaces. Note that also the classical result [17, Chapitre II, §4, No. 3, Corollaire 2] (see also Bonet, Pérez Carreras [25, Proposition 11.6.13]) of Grothendieck on the ultrabornologicity of $s' \hat{\otimes}_{\pi} s$ is a result on an ε -tensor product of a Fréchet space and a DF-space due to the nuclearity of s . Indeed, a certain nuclearity assumption is also the key point in results of Piśczek [26] (cf. also Domański [15]) on tensor products of PLS-spaces. As we explain at the end of Section 4, the latter results can be used in our situation; in fact [26, Theorem 6 and Theorem 9] provide criteria for tensor products of FS- and LS-spaces to be ultrabornological if at least one of them is nuclear.

We refer the reader to [8] for weighted spaces of continuous functions and to [19, 20, 21, 22, 25] for the general theory of locally convex spaces.

2. NOTATION AND PRELIMINARY RESULTS

Let X be a locally compact and σ -compact topological space. By $C(X)$ we denote the space of all continuous functions on X and by $C_c(X)$ the subspace of all functions with compact support. A weight on X is a strictly positive and continuous function on X . We consider a double sequence $\mathcal{A} = ((a_{N,n})_{n \in \mathbb{N}})_{N \in \mathbb{N}}$ of weights on X which is decreasing in n and increasing in N , i.e. $a_{N,n+1} \leq a_{N,n} \leq a_{N+1,n}$ holds for all N and n . This condition will be assumed on the double sequence $\mathcal{A}$ in the rest of this article. We define

$$C(a_{N,n})_0(X) := \{f \in C(X) ; a_{N,n}|f| \text{ vanishes at } \infty \text{ on } X\}.$$

Recall that a function $g: X \rightarrow \mathbb{R}$ is said to vanish at infinity on X if for each $\varepsilon > 0$ there is a compact set K in X such that $|g(x)| < \varepsilon$ for all $x \in X \setminus K$. The spaces $C(a_{N,n})_0(X)$ are Banach spaces for the norms $\|f\|_{N,n} := \sup_{x \in X} a_{N,n}(x)|f(x)|$, $f \in C(a_{N,n})_0(X)$. By the definition of these norms, $C(a_{N,n})_0(X) \subseteq C(a_{N,n+1})_0(X)$ holds with continuous inclusion for all N and n and we can define for each N the weighted inductive limit

$$(\mathcal{A}_N)_0 C(X) := \text{ind}_n C(a_{N,n})_0(X).$$

The latter space needs not to be regular. By Bierstedt, Meise, Summers [8, Theorem 2.6] it is regular if and only if it is complete and this is in turn equivalent to the fact that the sequence $\mathcal{A}_N := (a_{N,n})_{n \in \mathbb{N}}$ is regularly decreasing, see [8, Definition 2.1]. We refer to the latter article for more details on this condition. For each N we have $(\mathcal{A}_{N+1})_0 C(X) \subseteq (\mathcal{A}_N)_0 C(X)$ with continuous inclusion. Hence, $\mathcal{A}_0 C := ((\mathcal{A}_N)_0 C(X))_{N \in \mathbb{N}}$ is a projective spectrum of LB-spaces with inclusions as linking maps and we can form the following projective limit, called weighted PLB-space of continuous functions

$$(AC)_0(X) := \text{proj}_N (\mathcal{A}_N)_0 C(X),$$

which is the object of our study in this article. Since the space of continuous functions with compact support is contained in $(\mathcal{A}_N)_0 C(X)$ and since it is dense in each Banach space $C(a_{N,n})_0(X)$, it follows that the projective limit $(AC)_0(X)$, or rather the projective spectrum $\mathcal{A}_0 C$, is strongly reduced in the sense of

[38, Definition 3.3.5]. In fact it is a reduced projective limit in the sense of Köthe [20, p. 120], which is a stronger condition.

We refer the reader to the book of Wengenroth [38] for a detailed exposition of the theory of projective spectra of locally convex spaces $\mathcal{X} = (X_N)_{N \in \mathbb{N}}$, their projective limits $\text{proj}_N X_N$, the derived functor $\text{Proj}^1 \mathcal{X}$ and for conditions to ensure that the derived functor of a projective spectrum vanishes, i.e. that we have $\text{Proj}^1 \mathcal{X} = 0$, including important results by Palamodov [23, 24], Retakh [27], Braun, Vogt [13], Vogt [31, 33], Frerick, Wengenroth [16] and many others. At this point we only mention that, if $\mathcal{X} = (X_N)_{N \in \mathbb{N}}$ is a projective spectrum of locally convex spaces with inclusions as linking maps and limit $X = \text{proj}_N X_N$, the so-called fundamental resolution

$$0 \rightarrow X \rightarrow \prod_{N=1}^{\infty} X_N \xrightarrow{\sigma} \prod_{N=1}^{\infty} X_N,$$

where $\sigma((x_N)_{N \in \mathbb{N}}) := (x_{N+1} - x_N)$, is exact but σ is not necessarily surjective, what directs to the definition

$$\text{Proj}^1 \mathcal{X} := \left(\prod_{N=1}^{\infty} X_N \right) / \text{im } \sigma.$$

For more details see Wengenroth [38, Chapter 3].

In [34] Vogt introduced the condition (wQ). In the case of weighted PLB-spaces one can reformulate this condition in terms of the weights as follows. We say that the sequence $\mathcal{A}$ satisfies (wQ) if

$$\forall N \exists M \geq N, n \forall K \geq M, m \exists k, S > 0 : \frac{1}{a_{M,m}} \leq \max \left(\frac{S}{a_{N,n}}, \frac{S}{a_{K,k}} \right).$$

We have the following result concerning homological properties of the spectrum $\mathcal{A}_0 C$ and locally convex properties of $(AC)_0(X)$, which is due to Agethen, Bierstedt, Bonet [2].

Theorem 1. ([2, Theorem 3.7]) *The following are equivalent.*

- (i) $\text{Proj}^1 \mathcal{A}_0 C = 0$. (iii) $(AC)_0(X)$ is barrelled.
- (ii) $(AC)_0(X)$ is ultrabornological. (iv) $\mathcal{A}$ satisfies condition (wQ). □

Given a sequence of weights $\mathcal{A} = ((a_{N,n})_{n \in \mathbb{N}})_{N \in \mathbb{N}}$ we can also associate a weighted LF-space of continuous functions defined by

$$\mathcal{V}_0 C(X) := \text{ind}_n \text{proj}_N C(a_{N,n})_0(X).$$

Spaces of this type were investigated by Bierstedt, Bonet [6]. It is clear, that $\mathcal{V}_0 C(X) \subseteq (AC)_0(X)$ holds with continuous inclusion. The equality was investigated by Agethen, Bierstedt, Bonet [2] using the following condition introduced by Vogt [30, Satz 1.1]. We say that the sequence $\mathcal{A}$ satisfies condition (B) if

$$\forall (n(N))_{N \in \mathbb{N}} \subseteq \mathbb{N} \exists m \forall M \exists L, c > 0 : a_{M,m} \leq c \max_{N=1, \dots, L} a_{N, n(N)}.$$

Agethen, Bierstedt, Bonet [2] proved the following result on the forementioned equality of PLB- and LF-space.

Theorem 2. ([2, Theorem 3.10 and Corollary 3.11]) *The following statements hold.*

- (1) *If $\mathcal{A}$ satisfies condition (B), then $(AC)_0(X) = \mathcal{V}_0 C(X)$ holds algebraically. If, in addition, each $(\mathcal{A}_N)_0 C(X)$ is complete, then the converse also holds.*
- (2) *Assume that each $(\mathcal{A}_N)_0 C(X)$ is complete. Then $(AC)_0(X) = \mathcal{V}_0 C(X)$ holds algebraically and topologically if and only if $\mathcal{A}$ satisfies conditions (B) and (wQ).* □

In the sequel we complement the above by considering the special situation that the domain X is the product of two topological spaces X_1 and X_2 and the sequence $\mathcal{A}$ is the product of an increasing sequence $(a_N^1)_{N \in \mathbb{N}}$ defined on X_1 and a decreasing sequence $((a_n^2)^{-1})_{n \in \mathbb{N}}$ defined on X_2 . In this special setting we can associate a weighted Fréchet space to each of the sequences $(a_N^1)_{N \in \mathbb{N}}$ and $(a_n^2)_{n \in \mathbb{N}}$ and thus draw the line between properties of the PLB-space and the invariants (DN) and (Ω) for Fréchet spaces. In order to do this we have to characterize (DN) and (Ω) in terms of the weights.

3. (DN) AND (Ω) VS. (wQ)

Vogt [29] and Vogt, Wagner [35] introduced the following conditions, which are topological invariants of Fréchet spaces. We say that a Fréchet space E with a fundamental sequence of seminorms $(\|\cdot\|_n)_{n \in \mathbb{N}}$ satisfies condition (DN), if

$$\exists n \forall m \geq n, 0 < \theta < 1 \exists k \geq m, C > 0 \forall x \in E : \|x\|_m \leq C \|x\|_k^\theta \|x\|_n^{1-\theta}.$$

By Meise, Vogt [22, Lemma 29.10] the latter statement is equivalent to the original formulation [22, Definition on p. 359] of (DN). According to [22, Definition on p. 367] we say that a Fréchet space E with a fundamental sequence of seminorms $(\|\cdot\|_n)_{n \in \mathbb{N}}$ satisfies condition (Ω) , if

$$\forall N \exists M \geq N \forall K \geq M \exists D > 0, 0 < \theta < 1 \forall y \in E' : \|y\|_M^* \leq D(\|y\|_K^*)^\theta (\|y\|_N^*)^{1-\theta},$$

where $\|y\|_n^* := \sup_{\|x\|_n \leq 1} |y(x)|$. It is well-known (and an immediate consequence of [22, Lemma 29.13]) that replacing the above estimate by the inclusion $U_M \subseteq rU_N + Dr^{1-1/\theta}U_K$ (required for each $r > 0$ and with $U_n := \{x \in E : \|x\|_n \leq 1\}$ for $n \in \mathbb{N}$) yields an equivalent formulation of (Ω) .

Definition 3. Let $A = (a_n)_{n \in \mathbb{N}}$ be an increasing sequence of weights on a Hausdorff locally compact and σ -compact topological space X . We say that A satisfies condition $(DN)_w$ if

$$\exists n \forall m \geq n, 0 < \theta < 1 \exists k \geq m, C > 0 : a_m \leq Ca_k^\theta a_n^{1-\theta}.$$

We say that A satisfies condition $(\Omega)_w$ if

$$\forall N \exists M \geq N \forall K \geq M \exists D > 0, 0 < \theta < 1 : \frac{1}{a_M} \leq D\left(\frac{1}{a_K}\right)^\theta \left(\frac{1}{a_N}\right)^{1-\theta}.$$

Lemma 4. Let $A = (a_n)_{n \in \mathbb{N}}$ be an increasing sequence of weights on a Hausdorff locally compact and σ -compact topological space X . The Fréchet space $CA_0(X) = \text{proj}_n C(a_n)_0(X)$ satisfies condition (DN) if and only if the sequence A satisfies $(DN)_w$.

Proof. “ $\Leftarrow$ ” We choose n as in $(DN)_w$. For given $m \geq n, 0 < \theta < 1$ we choose $k \geq m$ and $C > 0$ as in $(DN)_w$. For an arbitrary $f \in CA_0(X)$ we obtain $\|f\|_m = \sup_{x \in X} a_m(x)|f(x)| \leq C \sup_{x \in X} a_k^\theta(x) a_n^{1-\theta}(x) |f(x)|^{\theta+1-\theta} \leq C \sup_{x \in X} (a_k(x)|f(x)|)^\theta \sup_{x \in X} (a_n(x)|f(x)|)^{1-\theta} = C \|f\|_k^\theta \|f\|_n^{1-\theta}$ and hence we have shown (DN).

“ $\Rightarrow$ ” Let $x_0 \in X$ be fixed. Since X is locally compact, there is a neighborhood filter $(K_\beta)_{\beta \in B}$ for x_0 consisting of compact sets. Since X is Hausdorff, $\cap_{\beta \in B} K_\beta = \{x_0\}$. For each K_β we choose a function $f_\beta \in C_c(X)$ with $f_\beta(x_0) = 1, 0 \leq f_\beta \leq 1$ and $\text{supp } f_\beta \subseteq K_\beta$. Thus we have $\cap_{\beta \in B} \text{supp } f_\beta = \{x_0\}$. Now we consider the net $(\sup_{x \in X} a(x)|f_\beta(x)|^\gamma)_{\beta \in B} \subseteq \mathbb{R}$ for some fixed weight a and $\gamma > 0$ and obtain $\sup_{x \in X} a(x)|f_\beta(x)|^\gamma = \sup_{x \in \text{supp } f_\beta} a(x)|f_\beta(x)|^\gamma \leq \sup_{x \in \text{supp } f_\beta} a(x) \rightarrow a(x_0)$ and $\sup_{x \in X} a(x)|f_\beta(x)|^\gamma \geq a(x_0)|f_\beta(x_0)|^\gamma = a(x_0)$.

Now we select n as in (DN). For given $m \geq n$ and $0 < \theta < 1$ we choose $k \geq m$ and $C > 0$ as in (DN). Now we fix some $x_0 \in X$ and consider the f_β defined above. We have $\sup_{x \in X} a_m(x)|f_\beta(x)| \leq C \sup_{x \in X} a_k^\theta(x)|f_\beta(x)|^\theta \sup_{x \in X} a_n^{1-\theta}(x)|f_\beta(x)|^{1-\theta}$ for each $\beta \in B$. Taking limits on each side yields the desired inequality, which finishes the proof since x_0 was arbitrary. $\square$

Lemma 5. Let $A = (a_n)_{n \in \mathbb{N}}$ be an increasing sequence of weights on a Hausdorff locally compact and σ -compact topological X . The Fréchet space $CA_0(X) = \text{proj}_n C(a_n)_0(X)$ satisfies condition (Ω) if and only if the sequence A satisfies $(\Omega)_w$.

Proof. “ $\Leftarrow$ ” We put $a := \frac{1}{a_N}$ and $b := \frac{1}{a_K}$ and use (cf. [22, proof of Lemma 29.13]) that $\min_{s>0}(sa + s^{1-1/\theta}b) = \theta^{-1}(\theta^{-1} - 1)^{\theta-1} a^{1-\theta} b^\theta$ holds for arbitrary a and $b > 0$. We put $C := \theta^{-1}(\theta^{-1} - 1)^{\theta-1} > 0$ and thus get $a^{1-\theta} b^\theta \leq \frac{1}{C}(sa + s^{1-1/\theta}b)$ for each $s > 0$. Hence we obtain $D\left(\frac{1}{a_K}\right)^\theta \left(\frac{1}{a_N}\right)^{1-\theta} \leq \frac{D}{C}\left(s\frac{1}{a_N} + s^{1-1/\theta}\frac{1}{a_K}\right) \leq \max\left(2\frac{D}{C}s\frac{1}{a_N}, 2\frac{D}{C}s^{1-1/\theta}\frac{1}{a_K}\right)$ for each $s > 0$. Now we define $r = 4\frac{D}{C}s$ and $D' = \frac{4D}{C}\left(\frac{C}{4D}\right)^{1-1/\theta} > 0$, hence $s = \frac{C}{4D}r$ and therefore $2\frac{D}{C}s^{1-1/\theta} = \frac{4D}{2C}\left(\frac{C}{4D}\right)^{1-1/\theta}r^{1-1/\theta} = \frac{D'}{2}r^{1-1/\theta}$. Replacing D' with D we get

$$(\star) \quad \forall N \exists M \geq N \forall K \geq M \exists D > 0, 0 < \theta < 1 \forall r > 0 : \frac{1}{a_M} \leq \max\left(\frac{r}{2a_N}, \frac{Dr^{1-1/\theta}}{2a_K}\right).$$

Now we show (Ω) in the second formulation mentioned at the beginning of this section. Let N be given. We choose $M \geq N$ as in $(\star)$. For given $K \geq M$ we select $D > 0$ and $0 < \theta < 1$ as in $(\star)$ and take an arbitrary $r > 0$. Let $f \in U_M$ be fixed, i.e. $|f| \leq \frac{1}{a_M} \leq \max\left(\frac{r}{2a_N}, \frac{Dr^{1-1/\theta}}{2a_K}\right)$. By [2, Lemma 3.4] there exist $\phi_1, \phi_2 \in C(X)$ with $0 \leq \phi_1, \phi_2 \leq 1, \phi_1 + \phi_2 = f$ such that $|\phi_1| \leq \frac{r}{a_N}, |\phi_2| \leq \frac{Dr^{1-1/\theta}}{a_K}$, i.e. $\phi_1 \in rU_N, \phi_2 \in Dr^{1-1/\theta}U_K$ and thus $f \in rU_N + Dr^{1-1/\theta}U_K$.

“ $\Rightarrow$ ” For a fixed N and for $x_0 \in X$ we consider $\delta_{x_0} : C(a_n)_0(X) \rightarrow \mathbb{C}, \delta_{x_0}(f) := f(x_0)$. Then we have $\|\delta_{x_0}\|_N^* = \sup_{f \in U_N} |\delta_{x_0}(f)| = \sup_{f \in U_N} |f(x_0)| \leq \frac{1}{a_N(x_0)}$. We choose $\varphi \in C_c(X)$ with $\varphi(x_0) = 1, 0 \leq \varphi \leq 1$ on X and put $f_0 := \frac{\varphi}{a_N}$. Then $f_0 \in C(a_N)_0(X)$ and $\sup_{x \in X} a_N(x)|f_0(x)| = \sup_{x \in X} a_N(x)\frac{\varphi(x)}{a_N(x)} = \sup_{x \in X} \varphi(x) =$

1, i.e. $f_0 \in U_N$. $\delta_{x_0}(f_0) = f_0(x_0) = \frac{\varphi(x_0)}{a_N(x_0)} = \frac{1}{a_N(x_0)}$ implies $\|\delta_{x_0}\|_N^* = \frac{1}{a_N(x_0)}$. $(\Omega)_w$ is the special case of (Ω) where we choose the functional to be δ_x for arbitrary $x \in X$. $\square$

After these preparations we are ready to investigate the situation we mentioned at the end of Section 2: For $i \in \{1, 2\}$ let X_i denote a locally compact and σ -compact Hausdorff topological space. Moreover, let $A^i = (a_n^i)_{n \in \mathbb{N}}$ be an increasing sequence of weights on X_i , i.e. $a_n^i \leq a_{n+1}^i$ for all $n \in \mathbb{N}$. We define the double sequence $\mathcal{A} = ((a_{N,n})_{N \in \mathbb{N}})_{n \in \mathbb{N}}$ by setting

$$a_{N,n} : X_1 \times X_2 \rightarrow \mathbb{R}, (x_1, x_2) \mapsto a_{N,n}(x_1, x_2) := [a_N^1 \otimes \frac{1}{a_n^2}](x_1, x_2) = \frac{a_N^1(x_1)}{a_n^2(x_2)}.$$

Thus, $\mathcal{A}$ satisfies the estimates $a_{N,n+1}(x_1, x_2) \leq a_{N,n}(x_1, x_2) \leq a_{N+1,n}(x_1, x_2)$ for all N, n and $(x_1, x_2) \in X_1 \times X_2$. In the sequel we refer to a sequence of the latter form by $\mathcal{A} = A^1 \otimes (A^2)^{-1}$. To simplify notation we put $X := X_1 \times X_2$ and consider the PLB-space $(AC)_0(X)$ in the notation established at the beginning of Section 2. In view of Theorem 1 we investigate if there is some relation between the conditions (DN) and (Ω) for the Fréchet spaces $C(A^i)_0(X_i)$ and the PLB-space $(AC)_0(X)$. According to the results above we can consider the weight conditions $(DN)_w$, $(\Omega)_w$ and (wQ) .

Proposition 6. *Let A^1 resp. A^2 be an increasing sequence of weights on X_1 resp. X_2 . Assume that A^1 satisfies $(\Omega)_w$ and that A^2 satisfies $(DN)_w$. Then the sequence $\mathcal{A} = A^1 \otimes (A^2)^{-1}$ on $X_1 \times X_2$ satisfies (wQ) .*

Proof. It suffices to show

$$\forall N \exists M \geq N, n \forall K \geq M, m \exists k, S > 0 \forall (x_1, x_2) \in X_1 \times X_2 : \frac{a_m^2(x_2)}{a_M^1(x_1)} \leq S \max\left(\frac{a_n^2(x_2)}{a_K^1(x_1)}, \frac{a_k^2(x_2)}{a_K^1(x_1)}\right).$$

In order to do this, let N be given. We select $M \geq N$ as in $(\Omega)_w$ and n as in $(DN)_w$. For given $K \geq M$ there exist $D > 0$ and $0 < \theta < 1$ with the estimate in $(\Omega)_w$. For arbitrary m and the same θ there exist $k \geq m$ and $C > 0$ with the estimate in $(DN)_w$. We put $S := 2CD$ and multiply the estimates in $(DN)_w$ and $(\Omega)_w$ to get $\frac{a_m^2(x_2)}{a_M^1(x_1)} \leq CD \left(\frac{a_k^2(x_2)}{a_K^1(x_1)}\right)^\theta \left(\frac{a_n^2(x_2)}{a_N^1(x_1)}\right)^{1-\theta} \leq CD \left(\frac{a_k^2(x_2)}{a_K^1(x_1)} + \frac{a_n^2(x_2)}{a_N^1(x_1)}\right) \leq S \max\left(\frac{a_k^2(x_2)}{a_K^1(x_1)}, \frac{a_n^2(x_2)}{a_N^1(x_1)}\right)$ for each $(x_1, x_2) \in X_1 \times X_2$. $\square$

In the light of Theorem 2 it is natural to ask whether there is any relation between the conditions (wQ) and (B); this was one of the objectives in [36]. In the latter work a variant of the following example was stated to show that “ $(wQ) \Rightarrow (B)$ ” is not true in general. Proposition 6 yields a very easy way of concluding that the double sequence $\mathcal{A}$ in Example 7 satisfies (wQ) ; we believe that this is less complicated than the approach of [36].

Example 7. In general, the assumptions of Proposition 6 do not imply that $\mathcal{A}$ satisfies condition (B). Let s be the Fréchet space of rapidly decreasing sequences. Then $s = \{x \in \mathbb{K}^{\mathbb{N}}; \forall k \in \mathbb{N} : \lim_{j \rightarrow \infty} j^k |x_j| = 0\} = CA_0(\mathbb{N})$ holds for $A = (a_k)_{k \in \mathbb{N}}$ with $a_k(j) = j^k$ by [22, Proposition 28.16]. In particular, s satisfies (DN) and (Ω) , see e.g. Meise, Vogt [22, Section III.29]. Now we consider $\mathcal{A} = ((a_N \otimes a_n^{-1})_{n \in \mathbb{N}})_{N \in \mathbb{N}}$ on $\mathbb{N} \times \mathbb{N}$, i.e. $(AC)_0(\mathbb{N} \times \mathbb{N}) = \text{proj}_N \text{ind}_n C(a_N \otimes a_n^{-1})_0(\mathbb{N} \times \mathbb{N})$, which satisfies (wQ) by Proposition 6. Straight forward calculations show that $\mathcal{A}$ does not satisfy (B).

Theorem 8. *Let A^1 resp. A^2 be an increasing sequence of weights on X_1 resp. X_2 . Consider the sequence $\mathcal{A} = A^1 \otimes (A^2)^{-1}$ on $X = X_1 \times X_2$ and assume that $C(A^1)_0(X_1)$ satisfies (Ω) and that $C(A^2)_0(X_2)$ satisfies (DN). Then $(AC)_0(X)$ is ultrabornological.*

Proof. The result follows immediately from Lemmas 4 and 5, Proposition 6 and Theorem 1. $\square$

Remark 9. Changing certain quantifiers in the definitions of (DN) and (Ω) , the latter turn into the weaker resp. stronger conditions $(\underline{DN})$ and $(\underline{\Omega})$. For the definition and recent results involving $(\underline{DN})$ and $(\underline{\Omega})$ we refer to Bonet, Domański [9, 10, 12].

Having $(\underline{DN})$ and $(\underline{\Omega})$ at hand it is easy to see how weight conditions $(\underline{DN})_w$ and $(\underline{\Omega})_w$ have to be defined in order to characterize the invariants $(\underline{DN})$ and $(\underline{\Omega})$ for weighted Fréchet spaces $CA_0(X)$. Then it is not hard to see that Proposition 6 and Theorem 8 are also valid with (DN) resp. (Ω) replaced by $(\underline{DN})$ resp. $(\underline{\Omega})$.

Since a detailed proof for these variants of Proposition 6 and Theorem 8 is straight forward, we omit it at this point; a detailed exposition can be found in [37]. However, the results might be of interest in view of applications and examples.

4. TENSOR PRODUCT REPRESENTATION

The constructions in the earlier sections already suggest the question, whether the space $(AC)_0(X)$ with an underlying sequence $\mathcal{A} = A^1 \otimes (A^2)^{-1}$ can be realized as the tensor product of a Fréchet and an LB-space. In the last section we deduce a representation of this kind, which finally will enable us (see Theorem 11) to utilize Theorem 8 to prove a criterion for the ultrabornologicity of an ε -tensor product of a weighted Fréchet space of continuous functions and a weighted LB-space of continuous functions. For further information on the general question of determining locally convex properties of tensor products of a Fréchet and a DF-space we refer to our comments in Section 1.

In what follows we need the so-called regularly decreasing condition; we refer to Bierstedt, Meise, Summers [8, Definition 2.1] for the definition and further information.

Proposition 10. *Let A^1 resp. A^2 be an increasing sequence of weights on X_1 resp. X_2 . Consider the double sequence $\mathcal{A} = A^1 \otimes (A^2)^{-1}$ on $X = X_1 \times X_2$ and assume that $\mathcal{V}^2 = (A^2)^{-1}$ is regularly decreasing. Then we have the isomorphism*

$$C(A^1)_0(X_1) \check{\otimes}_\varepsilon \mathcal{V}_0^2 C(X_2) \cong (AC)_0(X),$$

where $C(A^1)_0(X_1) = \text{proj}_N C(a_N^1)_0(X_1)$ resp. $\mathcal{V}_0^2 C(X_2) = \text{ind}_n C((a_n^2)^{-1})_0(X_2)$ is a weighted Fréchet resp. LB-space of continuous functions.

Proof. We compute

$$\begin{aligned} C(A^1)_0(X_1) \check{\otimes}_\varepsilon \mathcal{V}_0^2 C(X_2) &\stackrel{\text{dfn}}{=} (\text{proj}_N C(a_N^1)_0(X_1)) \check{\otimes}_\varepsilon (\text{ind}_n C((a_n^2)^{-1})_0(X_2)) \\ &\stackrel{(1)}{=} \text{proj}_N [C(a_N^1)_0(X_1) \check{\otimes}_\varepsilon \text{ind}_n C((a_n^2)^{-1})_0(X_2)] \\ &\stackrel{(2)}{=} \text{proj}_N \text{ind}_n [C(a_N^1)_0(X_1) \check{\otimes}_\varepsilon C((a_n^2)^{-1})_0(X_2)] \\ &\stackrel{(3)}{=} \text{proj}_N \text{ind}_n [C(a_N^1 \otimes (a_n^2)^{-1})_0(X_1 \times X_2)] \\ &\stackrel{\text{dfn}}{=} (AC)_0(X). \end{aligned}$$

The isomorphism (1) is true in general, see e.g. Jarchow [19, 16.3.2].

(2) can be seen as follows. We have

- a. [18, Theorem 4.1]: $C(a_N^1)_0(X_1)$ is an ε -space by Hollstein [18, Proposition 2.3],
- b. [18, Theorem 4.1.(ii)]: $\mathcal{V}_0^2 C(X_2)$ is quasi-complete, compact-regular (cf. Bierstedt, Meise, Summers [8, Corollary 2.7]) and all the $C((a_n^2)^{-1})_0(X_2)$ have the approximation property (cf. Bierstedt [3, Theorem 5.5.(3)]),
- c. [18, Proposition 4.4.(1)]: $C(a_N^1)_0(X_1)$ is Banach and $\mathcal{V}_0^2 C(X_2)$ is compact-regular (see b.),
- d. [18, Theorem 4.1., 2nd part]: $\mathcal{V}_0^2 C(X_2)$ and all the $C((a_n^2)^{-1})_0(X_2)$ are complete (see b.),

Therefore by Hollstein [18, Theorem 4.1] we have an isomorphism

$$\text{ind}_n [C(a_N^1)_0(X_1) \check{\otimes}_\varepsilon C((a_n^2)^{-1})_0(X_2)] \xrightarrow{\sim} C(a_N^1)_0(X_1) \check{\otimes}_\varepsilon \text{ind}_n C((a_n^2)^{-1})_0(X_2)$$

for each N . It is easy to see that the latter yields an isomorphism of the corresponding projective limits and we therefore get the desired isomorphism (2).

Finally, a result of Bierstedt [4, Theorem 1.2] yields an isomorphism

$$C(a_N^1)_0(X_1) \check{\otimes}_\varepsilon C((a_n^2)^{-1})_0(X_2) \xrightarrow{\sim} C(a_N^1 \otimes (a_n^2)^{-1})_0(X_1 \times X_2),$$

for each N and n which is given by $\sum_{i=1}^j f_i \otimes g_i \mapsto [(x_1, x_2) \mapsto \sum_{i=1}^j f_i(x_1)g_i(x_2)]$ – note that the equality $C(a_N^1)_0(X_1) \check{\otimes}_\varepsilon C((a_n^2)^{-1})_0(X_2) = C(a_N^1 \otimes (a_n^2)^{-1})_0(X_1 \times X_2)$ follows from Köthe [21, §44, 2.(5)], since both spaces are complete and $C((a_n^2)^{-1})_0(X_2)$ has the approximation property (see a. above). Again it is easy to see that the above isomorphisms yield an isomorphism of the corresponding PLB-spaces. This finally shows (3). $\square$

In the case of sequence spaces, the above tensor product is of the type $\lambda^0(A) \check{\otimes}_\varepsilon k^0(B)$, i.e. it is the tensor product of a Köthe echelon and Köthe coechelon space and thus Proposition 10 can be regarded as an extension of [2, Lemma 4.3] to continuous functions.

The space considered in Example 7, $(AC)_0(\mathbb{N} \times \mathbb{N})$ with $\mathcal{A} = ((a_N \otimes a_n^{-1})_{n \in \mathbb{N}})_{N \in \mathbb{N}}$ and $a_k(j) = j^k$, is by Proposition 10 isomorphic to the space $s \check{\otimes}_\varepsilon k^0(B)$ where $B = (j^{-k})_{j,k \in \mathbb{N}}$ (in this case the regularly

decreasing condition [7, Definition 3.1] can easily be verified). In fact, $(AC)_0(\mathbb{N} \times \mathbb{N})$ is isomorphic to $s \check{\otimes}_\varepsilon s'_b$ as Corollary 14 below will show.

Theorem 11. *Let A^1 resp. A^2 be an increasing sequence of weights on X_1 resp. X_2 . Assume that $\mathcal{V}^2 = (A^2)^{-1}$ is regularly decreasing, that $C(A^1)_0(X_1)$ satisfies (Ω) and that $C(A^2)_0(X_2)$ satisfies (DN) . Then the ε -tensor product $C(A^1)_0(X_1) \check{\otimes}_\varepsilon \mathcal{V}_0^2 C(X_2)$ of a Fréchet space and a DF-space is ultrabornological.*

Proof. The result follows directly from Theorem 8 and Proposition 10. $\square$

Remark 12. In view of Remark 9 it is clear that Theorem 11 is also valid if we replace (DN) resp. (Ω) by $(\overline{DN})$ resp. $(\overline{\Omega})$. Again we refer to [37] for details. $\square$

In the case of sequence spaces the meaning of Theorem 11 and Remark 12 is the following: Let $A = (a_n)_{n \in \mathbb{N}}$ be a Köthe matrix and $B = (b_n)_{n \in \mathbb{N}}$ be a decreasing sequence of strictly positive functions on $\mathbb{N}$ which is regularly decreasing. Put $B^{-1} = (b_n^{-1})_{n \in \mathbb{N}}$. Assume that $\lambda^0(A)$ satisfies (Ω) and $\lambda^0(B^{-1})$ satisfies (DN) or that $\lambda^0(A)$ satisfies $(\overline{\Omega})$ and $\lambda^0(B^{-1})$ satisfies $(\overline{DN})$. Then the two statements above imply that the space $\lambda^0(A) \check{\otimes}_\varepsilon k^0(B)$ is ultrabornological. The latter follows also from the results in [2, Section 4] and Proposition 6, since in [2, Section 4] ultrabornologicity of the space $\lambda^0(A) \check{\otimes}_\varepsilon k^0(B)$ was characterized via condition (wQ). Let us add that in the case of sequence spaces, B is regularly decreasing if and only if $\lambda^0(B^{-1})$ is quasnormable and that this is equivalent to condition (wS) of Bierstedt, Meise, Summers, see [7, Proposition on p. 48 and Proposition 3.2].

The setup of this section, in particular the fact that in Theorem 11 and Remark 12 one of the assumptions is an assumption on the space $C(A^2)_0(X_2)$ whereas the conclusion affects the space $\mathcal{V}_0^2 C(X_2)$ with $\mathcal{V}^2 = (A^2)^{-1}$, suggests the question if in the latter statements $\mathcal{V}_0^2 C(X_2)$ can be replaced by $C(A^2)_0(X_2)'_b$ or if dually $C(A^2)_0(X_2)$ can be replaced by $\mathcal{V}_0^2 C(X_2)'_b$.

For sequence spaces we have (in the notation above) $k^0(B)'_b \cong \lambda^1(B^{-1})$, see [5, Proposition 2.10]. If now $k^0(B)'_b$ satisfies (DN) , the same is true for $\lambda^1(B^{-1})$ and we can utilize [22, Exercise 29.8] in combination with arguments similar to the proof of [22, Lemma 29.10] to obtain that the Köthe matrix B^{-1} satisfies $(DN)_w$. Then Lemma 4 implies that $\lambda^0(B^{-1})$ satisfies (DN) . Thus, in the case of sequence spaces our second question can be answered positively as follows.

Corollary 13. *Let $A = (a_n)_{n \in \mathbb{N}}$ be a Köthe matrix and $B = (b_n)_{n \in \mathbb{N}}$ be a decreasing sequence of strictly positive functions on $\mathbb{N}$ which is regularly decreasing. Assume that $\lambda^0(A)$ satisfies (Ω) and $k^0(B)'_b$ satisfies (DN) or that $\lambda^0(A)$ satisfies $(\overline{\Omega})$ and $k^0(B)'_b$ satisfies $(\overline{DN})$. Then the space $\lambda^0(A) \check{\otimes}_\varepsilon k^0(B)$ is ultrabornological.* $\square$

In view of the second question, we have $\lambda^0(B^{-1})'_b \cong k^1(B)$, see [5, Proposition 2.10]. Moreover, if $\lambda^0(B^{-1})$ or equivalently $\lambda^0(B^{-1})'_b$ is nuclear then $k^1(B) \cong k^0(B)$ is valid (cf. [5, remark after Proposition 2.15]) and hence also our first question admits a positive answer in the case of (nuclear) sequence spaces. Let us add at this point that nuclearity of $\lambda^0(B^{-1})$ is equivalent to B satisfying the Grothendieck-Pietsch condition (N), see [7, Remark after Theorem 5.4], which implies the regularly decreasing condition by [7, Remark after Theorem 5.4 and Proposition 4.8].

Corollary 14. *Let $A = (a_n)_{n \in \mathbb{N}}$ and $B = (b_n)_{n \in \mathbb{N}}$ be Köthe matrices such that $\lambda^0(B)$, or equivalently its strong dual, is nuclear. Assume that $\lambda^0(A)$ satisfies (Ω) and $\lambda^0(B)$ satisfies (DN) or that $\lambda^0(A)$ satisfies $(\overline{\Omega})$ and $\lambda^0(B)$ satisfies $(\overline{DN})$. Then the space $\lambda^0(A) \check{\otimes}_\varepsilon \lambda^0(B)'_b$ is ultrabornological.* $\square$

In the general case of continuous functions, an analogon of Corollary 13 can be proved under the additional assumption that both factors of the tensor product are Schwartz and that one of them is nuclear – this is achieved by using the results [26, Theorem 6 and Theorem 9] of Piszczek (compare with Domański [15, Corollary 5.6]) on tensor products of PLS-spaces.

Proposition 15. *Let A be an increasing and $\mathcal{V}$ be a decreasing sequence of weights on X_1 resp. X_2 . Assume that the spaces $CA_0(X_1)$ and $\mathcal{V}_0 C(X_2)$ are Schwartz and that one of them is nuclear. If $CA_0(X_1)$ satisfies (Ω) and $\mathcal{V}_0 C(X_2)'_b$ satisfies (DN) or $CA_0(X_1)$ satisfies $(\overline{\Omega})$ and $\mathcal{V}_0 C(X_2)'_b$ satisfies $(\overline{DN})$, then the ε -tensor product $CA_0(X_1) \check{\otimes}_\varepsilon \mathcal{V}_0 C(X_2)$ is ultrabornological.*

Proof. Our general assumptions imply that $CA_0(X_1)$ and $\mathcal{V}_0C(X_2)$ both enjoy the dual interpolation estimate for big θ (in the case of (DN) and (Ω)) or small θ (in the case of $(\underline{DN})$ and $(\underline{\Omega})$), cf. [26, Proposition 1]. With [26, Theorem 6 resp. Theorem 9] our nuclearity assumptions imply that $CA_0(X_1) \hat{\otimes}_\varepsilon \mathcal{V}_0C(X_2)$ has the dual interpolation estimate for small or for big θ . Since $CA_0(X_1)$ and $\mathcal{V}_0C(X_2)$ are ultrabornological PLS-spaces, their completed ε -tensor product is a PLS-space, cf. [26, remarks at the end of Section 2]. Thus, $CA_0(X_1) \hat{\otimes}_\varepsilon \mathcal{V}_0C(X_2)$ is ultrabornological by Bonet, Domański [11, Corollary 1.2.(c)]. $\square$

In view of the arguments previous to Corollary 13, the sequence space case of Proposition 15 implies the sequence space cases of Theorem 11 and Remark 12, if we assume that $\lambda^0(A)$ and $\lambda^0(B^{-1})$ are Schwartz and that one of them is nuclear. In fact, the assumptions of Proposition 15 already imply $X_1 \cong X_2 \cong \mathbb{N}$ as the next statement shows.

Remark 16. Let X be a Hausdorff locally compact and σ -compact space. Let $A = (a_n)_{n \in \mathbb{N}}$ be an increasing sequence of weights on X and let $CA_0(X) = \text{proj}_n C(a_n)_0(X)$ be Schwartz. Then $X \cong \mathbb{N}$, where $\mathbb{N}$ is endowed with the discrete topology. The same conclusion is valid if we replace $CA_0(X)$ by $\mathcal{V}_0C(X) = \text{ind}_n C(v_n)_0(X)$ for a decreasing sequence $\mathcal{V} = (v_n)_{n \in \mathbb{N}}$ of weights.

Proof. Let $K \subseteq X$ be compact and consider the restriction map $q: CA_0(X) \rightarrow C(K)$. With the definition $C := \sup_{x \in K} 1/a_1(x)$ we have $a_1C \geq 1$ on K and thus $\sup_{x \in K} |f(x)| \leq C \sup_{x \in K} a_1(x) |f(x)| \leq C \sup_{x \in X} a_1(x) |f(x)|$ for each $f \in CA_0(X)$, i.e. q is continuous if we endow $C(K)$ with the sup-norm. Hence, q induces an isomorphism $CA_0(X)/\ker q \cong C(K)$ and thus $C(K)$ is a Schwartz space by [22, 24.18]. But $C(K)$ is also a Banach space, whence it is finite dimensional which in turn implies that K is finite (for infinite K it is not hard to construct an infinite set of continuous functions which is linearly independent); in particular, K has to carry the discrete topology. Now the conclusion follows since we assumed that X is σ -compact. In the case of $\mathcal{V}_0C(X)$ we may use the same arguments; the continuity of the restriction maps follows from the universal property of the inductive limit (e. g. [22, 24.7]). $\square$

To conclude, let us point out that Theorem 11 and Remark 12 clearly are more general in the case of weighted spaces of continuous functions whereas the results of Piśszczek and Domański which led us to Proposition 15 provide a general theory for (nuclear) PLS-space.

ACKNOWLEDGEMENTS. This article arises from a part of the author's doctoral thesis which was supervised by Klaus D. Bierstedt and José Bonet. The author thanks both of them for all their advice, helpful remarks and valuable suggestions. In addition, the author likes to thank the referee for the correction of an error in Lemma 5, for pointing out the relation of this work to the recent papers of Piśszczek and Domański and for his comments on the authors original proof of Proposition 6 (which was inspired by that of Vogt [32, Theorem 5.1] and involved slightly different but equivalent versions of $(DN)_w$ and $(\Omega)_w$), which led to a simplification of the presented results and in particular of the forementioned proof.

REFERENCES

- [1] Agethen, S. (2004). Spaces of continuous and holomorphic functions with growth conditions, Dissertation Universität Paderborn.
- [2] Agethen S., Bierstedt K. D., Bonet J. (2009). Projective limits of weighted (LB)-spaces of continuous functions, Arch. Math. (Basel) **92**, no. 5, 384–398.
- [3] Bierstedt K. D. (1973). Gewichtete Räume stetiger vektorwertiger Funktionen und das injektive Tensorprodukt. I, J. Reine Angew. Math. **259**, 186–210.
- [4] Bierstedt K. D. (1974). Injektive Tensorprodukte und Slice-Produkte gewichteter Räume stetiger Funktionen, J. Reine Angew. Math. **266**, 121–131.
- [5] Bierstedt K. D. (1988). An introduction to locally convex inductive limits, Functional Analysis and its Applications (Nice, 1986) 35–133, ICPAM Lecture Notes, World Sci. Publishing, Singapore, 35–133.
- [6] Bierstedt K. D., Bonet J. (1994). Weighted (LF)-spaces of continuous functions, Math. Nachr. **165**, 25–48.
- [7] Bierstedt K. D., Meise R., Summers W. H. (1982). Köthe sets and Köthe sequence spaces, pp. 27–91 in *Functional analysis, holomorphy and approximation theory (Rio de Janeiro, 1980)*, North-Holland Math. Stud. **71**.
- [8] Bierstedt K. D., Meise R., Summers W. H. (1982) A projective description of weighted inductive limits, Trans. Amer. Math. Soc. **272**, no. 1, 107–160.
- [9] Bonet, J., Domański, P. (2001). Parameter dependence of solutions of partial differential equations in spaces of real analytic functions, Proc. Amer. Math. Soc. **129**, no. 2, 495–503.
- [10] Bonet, J., Domański, P. (2006). Parameter dependence of solutions of differential equations on spaces of distributions and the splitting of short exact sequences, J. Funct. Anal. **230**, no. 2, 329–381.
- [11] Bonet, J., Domański, P. (2007). The structure of spaces of quasianalytic functions of Romieu type, Arch. Math. (Basel) **89**, no. 5, 430–441.
- [12] Bonet, J., Domański, P. (2008). The splitting of exact sequences of PLS-spaces and smooth dependence of solutions of linear partial differential equations, Adv. Math. **217**, no. 2, 561–585.

- [13] Braun, R., Vogt, D. (1997). A sufficient condition for $\text{Proj}^1 \mathcal{X} = 0$, *Michigan Math. J.* **44**, no. 1, 149–156.
- [14] Domański, P. (2004). Classical PLS-spaces: spaces of distributions, real analytic functions and their relatives, *Orlicz centenary volume*, Banach Center Publ. **64**, 51–70.
- [15] Domański, P. (2010). Real analytic parameter dependence of solutions of differential equations, *Rev. Mat. Iberoamericana* **26**, 174–238.
- [16] Frerick, L., Wengenroth, J. (1996). A sufficient condition for vanishing of the derived projective limit functor, *Arch. Math. (Basel)* **67**, no. 4, 296–301.
- [17] Grothendieck, A. (1955). *Produits Tensoriels Topologiques et Espaces Nucléaires*, *Mem. Am. Math. Soc.* **16**.
- [18] Hollstein, R. (1980). Inductive limits and ε -tensor products, *J. Reine Angew. Math.* **319**, 38–62.
- [19] Jarchow, H. (1981). *Locally Convex Spaces*, B. G. Teubner, Stuttgart.
- [20] Köthe, G. (1969). *Topological vector spaces. I*, Grundlehren der math. Wiss. **159**, Springer-Verlag, New York.
- [21] Köthe, G. (1979). *Topological vector spaces. II*, Grundlehren der math. Wiss. **237**, Springer-Verlag, New York.
- [22] Meise, R., Vogt, D. (1997). *Introduction to Functional Analysis*, Oxford Graduate Texts in Math. **2**, The Clarendon Press, Oxford University Press, New York.
- [23] Palamodov, V. P. (1968). The projective limit functor in the category of topological linear spaces, *Mat. Sb.* **75**, 567–603 (in Russian), English transl. *Math. USSR Sbornik* **17**, 189–315.
- [24] Palamodov, V. P. (1971). Homological methods in the theory of locally convex spaces, *Uspekhi Mat. Nauk* **26**(1), 3–66 (in Russian), English transl., *Russian Math. Surveys* **26**(1), 1–64.
- [25] Pérez Carreras, P., Bonet, J. (1987). *Barrelled Locally Convex Spaces*, North-Holland Mathematics Studies **113**.
- [26] Piszczek, K. (2010). On a property of PLS-spaces inherited by their tensor products, *Bull. Belg. Math. Soc. Simon Stevin* **17**, 155–170.
- [27] Retakh, V. S. (1971). Subspaces of a countable inductive limit (English transl.), *Soviet Math. Dokl.* **11**, 1384–1386.
- [28] Varol, O. (2007). A generalization of a theorem of A. Grothendieck, *Math. Nachr.* **280**, no. 3, 313–325.
- [29] Vogt, D. (1977). Charakterisierung der Unterräume von s , *Math. Z.* **155**, no. 2, 109–117.
- [30] Vogt, D. (1983). Frécheträume zwischen denen jede stetige lineare Abbildung beschränkt ist, *J. Reine Angew. Math.* **345**, 182–200.
- [31] Vogt, D. (1987). Lectures on projective spectra of (DF)-spaces, Seminar lectures, Wuppertal.
- [32] Vogt, D. (1987). On the functors $\text{Ext}^1(E, F)$ for Fréchet spaces, *Studia Math.* **85**, no. 2, 163–197.
- [33] Vogt, D. (1989). Topics on projective spectra of (LB)-spaces, *Adv. in the Theory of Fréchet spaces (Istanbul, 1988)*, NATO Adv. Sci. Inst. Ser. C **287**, 11–27.
- [34] Vogt, D. (1992). Regularity properties of (LF)-spaces, *Progress in Functional Analysis, Proc. Int. Meet. Occas. 60th Birthd. M. Valdivia*, Peñáscola/Spain, North-Holland Math. Stud. **170**, 57–84.
- [35] Vogt, D., Wagner, M.-J. (1980). Charakterisierung der Quotientenräume von s und eine Vermutung von Martineau, *Studia Math.* **67**, no. 3, 225–240.
- [36] Wegner, S.-A. (2007). Inductive kernels and projective hulls for weighted (PLB)- and (LF)-spaces of continuous functions, Diplomarbeit Universität Paderborn.
- [37] Wegner, S.-A. (2010). Projective limits of weighted (LB)-spaces of holomorphic functions, PhD-thesis, Universidad Politécnica de Valencia.
- [38] Wengenroth, J. (2003). *Derived Functors in Functional Analysis*, Lecture Notes in Mathematics **1810**, Springer, Berlin.